

CTGATGGTGGCCAAATGGACAGGGGACTTTTTCAATAAGGGCATTATGATATCCACGTGGGCCTGCGAGC
TCTGGAATGGGAGACAGAGGTGGAAATGGACAAGCTGAGAGCCAGCGACATCATGGAGCCCAACCTGAC
CGCACACCCGCATCCAGTCTCTGGTGAGCATCCTGCGCACCACGGTCCACCATGCCTTCCCAGGTGGTCACA
GGTAACGAGAAGGAGTTCATGAAGGGCAACCAGCTCATCAGCAACAACATCAAGTTCAAGAAATCCAGCA
GGCTGGCGAGCAGCGCAAACGGAGCCAGTCCATGAAGTCCTACCCATCCAGCGAGCTACGGAACATGTGT
TCGCCTCTGAGGAGCCAGCCGAGAAGGAGGACCTCCTGCAGCAGATGCTGGAAAGGAGATACTCCCTA
TACCCTGACCAGTCCCCAAGTGAAGACTGGACCATGGAGGAGCGGTTCCGCCCTCTGACCTTCCACGGCCT
GTCGCAGCTTGTCACCCTGCTTGTCCGAGGAGTTTGTACTCTGAAAGCCAGTCGAGCGCCAGCCAGCCGO
ATGCCGAGATGGCCGAGGACTACCCGCGGTACCCCGACATCCACGACCTGGACCTGACGCTGCTCAACCC
GTGGATGTCACCCCATACATGAACCCTTCGCCTTTCACCGTCTCGCCCAACACCCACGTCTCCAAGTCTTC
CAGAACGATGGGCCTGCGCCACCTGCCCGTGGTGAACGCTGTGGGAGAGATCGTGGGGATCATCACACGG
CCTATGAATTTCTGCAGGCCCGGCTGAGGCAGCACTACCAGACCATCTTGCCAACCTTCTTGACAAAGTG
CCTATCCCTAACCTCTCCTCGGTCTCGATTCTACGTAG

Parent Cell Line: HEK293

Quantity: 1 vial of ~1 x 10⁶ cells; froze

Thawing Procedure:

The cells must be thawed in a 37 °C water bath under 2 min. Using aseptic technique, add the contents of the vial to 9 ml of complete growth medium, centrifuge for 5 min at 125 x g. Aspirate the medium, and resuspend in 10 mL of complete growth medium, and place into a culture vessel of your choice. Only add selection antibiotic to the medium after 24 hours in culture.